

Light-cone divergence in twist-3 correlation functions

L.P. Gamberg^a, D.S. Hwang^b, A. Metz^c, and M. Schlegel^c

^a*Division of Science, Penn State Berks, Reading, PA 19610, USA*

^b*Department of Physics, Sejong University, Seoul 143-747, Korea*

^c*Institut für Theoretische Physik II,
Ruhr-Universität Bochum, D-44780 Bochum, Germany*

February 7, 2008

Abstract

It is argued that the definition of the twist-3 transverse momentum dependent correlation functions must be modified if they contain light-like Wilson lines. In the framework of a simple spectator model of the nucleon we show explicitly the presence of a light-cone divergence for a specific twist-3 time-reversal odd parton density. This divergence emerges for all eight twist-3 T-odd correlators and appears also in the case of a quark target in perturbative QCD. The divergence can be removed by using non-light-like Wilson lines. Based on our results we argue that currently there exists no established factorization formula for transverse momentum dependent twist-3 observables in semi-inclusive DIS and related processes.

Partonic correlation functions (distributions and fragmentation functions) that depend on both the longitudinal and transverse parton momentum (p_T -dependent) have received considerable attention over the past years. The most prominent examples are the (naive) time-reversal odd (T-odd) Sivers parton density [1, 2] and the Collins fragmentation function [3], where the status of the mere existence of T-odd parton distributions was clarified only relatively recently [4, 5]. A great deal of theoretical and experimental effort has been devoted to determining these functions by characterizing and measuring single spin asymmetries in semi-inclusive deep-inelastic scattering (DIS) [6, 7, 8] and azimuthal asymmetries in e^+e^- -annihilation [9] (see also [10, 11]).

The first experimental studies on single spin asymmetries in semi-inclusive DIS dealt with the observables A_{UL} (unpolarized lepton beam and longitudinally polarized target) [12, 13, 14, 15] and A_{LU} (longitudinally polarized lepton beam and unpolarized target) [16]. By contrast they are suppressed like $1/Q$ (with Q denoting the virtuality of the exchanged photon) with respect to the leading twist asymmetry A_{UT} (transversely polarized target). Consequently, a parton model description of these asymmetries invokes (p_T -dependent) twist-3 correlation functions [17, 18, 19] (see, e.g., Refs. [20, 21, 22, 23, 24, 25, 26, 27, 28] for related theoretical work on twist-3 single spin asymmetries). A complete list of p_T -dependent twist-3 correlation functions was obtained only recently [29, 30, 31, 32, 33].

It is non-trivial to establish QCD-factorization for transverse momentum dependent hard scattering processes in terms of p_T -dependent partonic correlators. While the twist-2 case was studied in some detail in the literature (see, e.g., Refs. [34, 35, 36, 37, 38]), for twist-3 observables only a tree-level formalism is available up to now [17, 18, 19, 39]. To some extent the difficulty is intimately related to the task of finding a suitable Wilson line that ensures color gauge invariance of the p_T -dependent correlation functions (see in particular the discussion in [40]). In this context

it was emphasized already in Ref. [41], that light-like Wilson lines typically lead to divergences. At low-order in α_{qcd} , in calculations of twist-2 p_T -dependent correlators with light-like Wilson lines indeed certain diagrams show such a light-cone divergence. On the other hand, one-loop computations of the T-odd Sivers function defined with a light-like Wilson line yield well-behaved results [5, 42, 43, 44, 45, 46].

The main intention of this note is to delineate the presence of a light-cone divergence explicit in the case of twist-3 T-odd correlators with light-like Wilson lines. To this end we perform a one-loop calculation of the particular twist-3 T-odd parton distribution $g^\perp$ [32, 33] which appears in the description of the beam spin asymmetry A_{LU} [32]. Our study is based on a simple diquark spectator model of the nucleon. The light-cone divergence which appears for $g^\perp$ also shows up for all twist-3 T-odd parton distributions. Moreover, one arrives at the same conclusion when using a quark target in perturbative QCD. It is therefore evident that twist-3 p_T -dependent correlators containing light-like Wilson lines are undefined. The presence of the light-cone divergence also implies that the frequently used tree-level formalism of Refs. [17, 18, 19, 39] for twist-3 observables in semi-inclusive DIS and related processes at least needs a modification. Moreover, the so-called subtraction formalism [48, 36, 37, 38] used for an all-order description of p_T -dependent twist-2 processes cannot directly be applied in the twist-3 case. As a consequence, currently there exists no established factorization formula for processes like the longitudinal twist-3 single spin asymmetries A_{LU} and A_{UL} .

Now, we discuss elements of the calculation of $g^\perp$. The starting point is the quark-quark correlation function

$$\begin{aligned} \Phi_{ij}^{(v)}(x, \vec{p}_T; P, S) = & \int \frac{d\xi^- d^2\vec{\xi}_T}{(2\pi)^3} e^{ip\cdot\xi} \langle P, S | \bar{\Psi}_j(0) \mathcal{P} \exp \left\{ -ig \int_0^\infty d\lambda (v \cdot A)(\lambda v) \right\} \\ & \times \mathcal{P} \exp \left\{ ig \int_0^\infty d\lambda (v \cdot A)(\lambda v + \xi) \right\} \Psi_i(\xi) | P, S \rangle \Big|_{\xi^+=0}, \quad (1) \end{aligned}$$

where P , S , and p respectively denote the nucleon momentum, nucleon spin, and quark momentum. We choose a frame in which the light-cone momentum P^+ is large, and the plus-component of the quark momentum is given by $p^+ = xP^+$. The two Wilson lines that appear in the definition (1) are determined by the vector $v = (v^+, v^-, \vec{0}_T)$, with the proper light-like limit given by $(0, 1, \vec{0}_T)$. Note that the twist-3 formalism of Refs. [17, 18, 19, 39] is working with light-like lines. The Wilson lines are of particular importance to enforce color gauge invariance in the quark correlation functions as well as to provide a mechanism to generate final state interactions and the associated complex phases necessary to characterize the T-odd parton densities [4, 5].

We perform a one-loop calculation in Feynman gauge for which a transverse gauge link at light-cone infinity [42, 47] is not relevant. The function $g^\perp$ is defined via the correlator in (1) by means of [32, 33]

$$2\varepsilon_T^{ij} p_{Tj} g^\perp(x, \vec{p}_T^2) = -P^+ \text{Tr} \left(\Phi(x, \vec{p}_T) \gamma^i \gamma_5 \right), \quad (2)$$

where the conventions $\varepsilon^{0123} = 1$ and $\varepsilon_T^{ij} \equiv \varepsilon^{-+ij}$ are used. Note that in (2) we take an average over the nucleon polarization, as the function $g^\perp$ is defined for an unpolarized target.

In the calculation the vector v is taken to be slightly off the light-cone ($v^+ \neq 0$, $|v^+|/v^- \ll 1$) and space-like ($v^2 < 0$). For our purpose here one equally well could use a time-like Wilson line. The light-cone divergence of $g^\perp$ will show up in the light-like limit $|v^+|/v^- \rightarrow 0$. Non-light-like Wilson lines as regulators were already used much earlier in connection with a proper definition of p_T -dependent correlation functions [41].




Figure 1: One-loop diagram relevant to the calculation of $g^\perp$ for the quark in the scalar diquark model of Ref. [4]. [h.c.] stands for the Hermitian Conjugate diagram. The thin dashed line depicts the cut of the correlator. In this diagram the diquark (dashed line) interacts through the exchange of a photon with the eikonalized quark (double line).

We compute the correlator in the simple diquark spectator model of the nucleon used in Ref. [4]. In this approach the interaction between the nucleon, the quark, and the diquark is given by a point-like scalar vertex with the coupling constant λ . We use the model with an Abelian gauge boson. In order to determine the distribution $g^\perp$ to lowest non-trivial order the diagram in Fig. 1 has to be evaluated, where only the imaginary part of the loop amplitude (with loop-momentum l) is relevant. The contribution of this diagram to the correlator is given by

$$\begin{aligned} \Phi_{ij}^{(v)}(x, \vec{p}_T; P, S) &= i \frac{e_q e_s \lambda^2}{2(2\pi)^3} \frac{1}{(\vec{p}_T^2 + \tilde{m}^2) P^+} \int \frac{d^4 l}{(2\pi)^4} v \cdot (2P - 2p + l) \\ &\times \left\{ \frac{[\bar{u}(P, S)(\not{p} - \not{l} + m_q)]_j [(\not{p} + m_q)u(P, S)]_i}{[(l \cdot v) + i\varepsilon][l^2 - i\varepsilon][(p - l)^2 - m_q^2 - i\varepsilon][(P - p + l)^2 - m_s^2 - i\varepsilon]} \right. \\ &\left. - \frac{[\bar{u}(P, S)(\not{p} + m_q)]_j [(\not{p} - \not{l} + m_q)u(P, S)]_i}{[(l \cdot v) - i\varepsilon][l^2 + i\varepsilon][(p - l)^2 - m_q^2 + i\varepsilon][(P - p + l)^2 - m_s^2 + i\varepsilon]} \right\}, \quad (3) \end{aligned}$$

with $\tilde{m}^2 \equiv x(1-x) \left(-M^2 + \frac{m_q^2}{x} + \frac{m_s^2}{1-x} \right)$ and quark momentum in light-cone coordinates given by $p = \left(xP^+, -\frac{\vec{p}_T^2 + m_s^2 - M^2(1-x)}{2P^+(1-x)}, \vec{p}_T \right)$. Using the definition in Eq. (2) the calculation of $g^\perp$ is straightforward and one arrives at the expression

$$\begin{aligned} \varepsilon_T^{ij} p_{Tj} g^\perp(x, \vec{p}_T^2) &= -\frac{e_q e_s \lambda^2}{2(2\pi)^3} \frac{1}{(\vec{p}_T^2 + \tilde{m}^2)} \sum_{\pm} \int \frac{d^4 l}{(2\pi)^4} \\ &\times \frac{v \cdot (2P - 2p + l) \left[\varepsilon_T^{ij} p_{Tj} (P^+ l^- - P^- l^+) + \varepsilon_T^{ij} l_{Tj} (P^- p^+ - P^+ p^-) \right]}{[(l \cdot v) \pm i\varepsilon][l^2 \mp i\varepsilon][(p - l)^2 - m_q^2 \mp i\varepsilon][(P - p + l)^2 - m_s^2 \mp i\varepsilon]}. \quad (4) \end{aligned}$$

The difficulty with the use of light-like lines can be understood by considering the dependence on the loop integration variable l in Eq. (4). If the vector v is taken to be light-like the l^- -dependence in the propagator that is due to the Wilson line drops out. As a consequence the integral diverges (logarithmically) for $l^+ \rightarrow 0$ and $l^- \rightarrow \pm\infty$. The crucial point is the presence of the factor l^- in the numerator of (4). This is in contrast to the corresponding calculations of the twist-2 T-odd Sivers function $f_{1T}^\perp$ and Boer-Mulders function $h_1^\perp$, for which the box graph in Fig. 1 provides a well-defined finite result when using light-like Wilson lines. We also note that, as far as a light-cone

divergence is concerned, the contribution of the box graph to the six twist-2 T-even p_T -dependent parton densities is well behaved in the light-like limit, because there appears no factor of l^- in the numerator. Here we do not address possible complications (for T-even functions) in the case of a massless gluon.

Performing the loop-integral in (4) yields the result

$$g^\perp(x, \vec{p}_T^2) = \frac{e_q e_s \lambda^2}{4(2\pi)^4} \frac{1}{(\vec{p}_T^2 + \tilde{m}^2)} \left\{ \frac{1-x}{x} \ln \left(\frac{|v^+| \tilde{m}^2}{v^-(P^+)^2} \right) - \frac{1-x}{x} \left(\ln(2x(1-x)) - \frac{2}{2-x} \ln \left(\frac{x}{2} \right) \right) \right. \\ \left. + \frac{(1-x)\vec{p}_T^2 + x m_s^2 - x(1-x)^2 M^2}{x \vec{p}_T^2} \ln \left(\frac{\vec{p}_T^2 + \tilde{m}^2}{\tilde{m}^2} \right) \right\} + \mathcal{O} \left(\frac{|v^+|}{v^-} \right), \quad (5)$$

where we made an expansion in the (small) parameter $|v^+|/v^-$ in order to simplify the algebra. The first term in the brackets in (5) makes the logarithmic light-cone divergence explicit. Note also that the result is boost-invariant and that, due to the expansion in $|v^+|/v^-$, the result cannot be used in the limit $x \rightarrow 0$. Eventually, we mention that in the case of the corresponding treatment of the Sivers function with a non-light-like line we recover the well-known (finite) result [5, 42, 44, 45, 46] in the light-like limit by putting $|v^+|/v^- \rightarrow 0$ at the end of the calculation.

Let us also point out that the light-cone divergence can be avoided by means of a phenomenological approach in the sense that we replace the proton-quark-diquark coupling according to

$$\lambda \rightarrow \lambda \frac{1}{\left(1 - \frac{p^2}{\Lambda^2} - i\varepsilon\right)}, \quad (6)$$

i.e., we introduce a monopole form factor by hand. In Eq. (6), p denotes the quark momentum while Λ is an arbitrary mass scale. Then the expression in (4) is replaced by

$$\varepsilon_T^{ij} p_{Tj} g^\perp(x, \vec{p}_T^2) = \frac{e_q e_s \lambda^2}{2(2\pi)^3} \frac{\Lambda^4(1-x)}{(\vec{p}_T^2 + \tilde{m}^2)(\vec{p}_T^2 + \tilde{m}_\Lambda^2)} \sum_{\pm} \int \frac{d^4 l}{(2\pi)^4} \\ \times \frac{(2P^+(1-x) + l^+) [\varepsilon_T^{ij} p_{Tj} (P^+ l^- - P^- l^+) + \varepsilon_T^{ij} l_{Tj} (P^- p^+ - P^+ p^-)]}{[l^+ \pm i\varepsilon][l^2 \mp i\varepsilon][(p-l)^2 - m_q^2 \mp i\varepsilon][(P-p+l)^2 - m_s^2 \mp i\varepsilon][(p-l)^2 - \Lambda^2 \mp i\varepsilon]}, \quad (7) \\ \text{with } \tilde{m}_\Lambda^2 \equiv x(1-x) \left(-M^2 + \frac{\Lambda^2}{x} + \frac{m_s^2}{1-x} \right).$$

In (7) we use the light-like Wilson line because using this form factor results in an additional factor of l^- in the denominator rendering the integral finite in the potentially dangerous region $l^+ \rightarrow 0$ and $l^- \rightarrow \pm\infty$. Although we get a finite result in this way we want to emphasize again that such an approach is purely phenomenological and cannot be derived using a Lagrangian of a microscopic theory/model.

We end this short note by summarizing the main points and providing some additional discussion.

- We have shown that twist-3 T-odd p_T -dependent parton densities cannot be defined by using light-like Wilson lines. In order to make this point explicit we have presented a one-loop calculation of the parton distribution $g^\perp$, which appears for instance in the beam spin asymmetry A_{LU} in semi-inclusive DIS. We arrive at the same conclusion for all eight twist-3 T-odd parton densities of a spin-1/2 particle for which we pushed the calculation up to the level of Eq. (4).

- While our calculation is based on a simple diquark spectator model of the nucleon, we find the same light-cone divergence also in the case of a quark target in perturbative QCD. Therefore, the main conclusion we have drawn is not based on some artefact of a simple model.
- With regard to possible light-cone divergences from Fig. 1, all eight twist-2 p_T -dependent parton densities are well-behaved in the light-like limit.
- Although we expect that the same divergence problem arises from the graph in Fig. 1 for T-even twist-3 parton densities, at present we cannot exclude that there appears a cancellation of divergent terms once all one-loop diagrams are taken into account.
- For the T-odd functions we considered here the light-cone divergence can be regularized by using non-light-like Wilson lines (see, e.g., also Ref. [41]). We introduced such lines by hand. It would be very interesting to investigate (as a first step) if a tree-level factorization formalism in the spirit of Refs. [17, 18, 19, 39] can be established which automatically provides definitions of twist-3 p_T -dependent correlators with non-light-like Wilson lines.
- In the literature a subtraction method has been used (see, e.g., Refs. [48, 36, 37, 38]) in order to formulate all-order factorization for transverse momentum dependent twist-2 observables. In this formalism also correlation functions with non-light-like lines appear. However, in particular when performing one-loop calculations of an observable, the dependence on various non-light-like directions drops out by construction, as it should be. In contrast, using this formalism in the case of, e.g., the longitudinal asymmetry A_{LU}^{jet} for jet production in DIS, which exclusively is given by $g^\perp$ [32], leads to a result depending on the a priori arbitrary ratio $|v^+|/v^-$. It is quite possible that the subtraction formalism can be generalized such that it can cope with transverse momentum dependent twist-3 observables. But this requires further studies.
- In a very recent manuscript [49] also a result for $g^\perp$ in the scalar diquark model has been provided. Rather than starting from an operator definition, this result is extracted from the calculation of the beam spin asymmetry A_{LU} in semi-inclusive DIS. By doing so and assuming factorization in the spirit of Refs. [17, 18, 19, 39] one arrives at an expression for $g^\perp$ which does not show the divergence discussed in our note, because the result for the full asymmetry A_{LU} is finite in the diquark spectator model [30, 31, 49]. However, apparently the factorization assumption is not valid for this twist-3 process, and $g^\perp$ obtained in Ref. [49] is not the corresponding twist-3 distribution function defined through Eqs. (1) and (2).

Acknowledgements: We thank Gary R. Goldstein for useful discussions. The work of M.S. has been supported by the Graduiertenkolleg “Physik der Elementarteilchen an Beschleunigern und im Universum.” The work has also been partially supported by the KOSEF (Korea Science and Engineering Foundation), by the Verbundforschung (BMBF), and by the Transregio/SFB Bochum-Bonn-Giessen. This research is part of the EU Integrated Infrastructure Initiative Hadronphysics Project under contract number RII3-CT-2004-506078.

References

- [1] D.W. Sivers, Phys. Rev. D **41**, 83 (1990).
- [2] D.W. Sivers, Phys. Rev. D **43**, 261 (1991).
- [3] J.C. Collins, Nucl. Phys. **B396**, 161 (1993).

- [4] S.J. Brodsky, D.S. Hwang, and I. Schmidt, Phys. Lett. B **530**, 99 (2002).
- [5] J.C. Collins, Phys. Lett. B **536**, 43 (2002).
- [6] A. Airapetian *et al.* [HERMES Collaboration], Phys. Rev. Lett. **94**, 012002 (2005).
- [7] V.Y. Alexakhin *et al.* [COMPASS Collaboration], Phys. Rev. Lett. **94**, 202002 (2005).
- [8] M. Diefenthaler, arXiv:hep-ex/0507013.
- [9] R. Seidl *et al.* [Belle Collaboration], Phys. Rev. Lett. **96**, 232002 (2006).
- [10] W. Vogelsang and F. Yuan, Phys. Rev. D **72**, 054028 (2005).
- [11] A.V. Efremov, K. Goeke, and P. Schweitzer, Phys. Rev. D **73**, 094025 (2006).
- [12] A. Airapetian *et al.* [HERMES Collaboration], Phys. Rev. Lett. **84**, 4047 (2000).
- [13] A. Airapetian *et al.* [HERMES Collaboration], Phys. Rev. D **64**, 097101 (2001).
- [14] A. Airapetian *et al.* [HERMES Collaboration], Phys. Lett. B **562**, 182 (2003).
- [15] A. Airapetian *et al.* [HERMES Collaboration], Phys. Lett. B **622**, 14 (2005).
- [16] H. Avakian *et al.* [CLAS Collaboration], Phys. Rev. D **69**, 112004 (2004).
- [17] J. Levelt and P.J. Mulders, Phys. Lett. B **338**, 357 (1994).
- [18] P.J. Mulders and R.D. Tangerman, Nucl. Phys. **B461**, 197 (1996) [Erratum-ibid. **B484**, 538 (1997)].
- [19] D. Boer and P.J. Mulders, Phys. Rev. D **57**, 5780 (1998).
- [20] K.A. Oganessian, H.R. Avakian, N. Bianchi, and A.M. Kotzinian, arXiv:hep-ph/9808368.
- [21] A.V. Efremov, K. Goeke, M.V. Polyakov, and D. Urbano, Phys. Lett. B **478**, 94 (2000).
- [22] E. De Sanctis, W.D. Nowak, and K.A. Oganessian, Phys. Lett. B **483**, 69 (2000).
- [23] B.Q. Ma, I. Schmidt, and J.J. Yang, Phys. Rev. D **63**, 037501 (2001).
- [24] A.V. Efremov, K. Goeke, and P. Schweitzer, Phys. Lett. B **522**, 37 (2001) [Erratum-ibid. B **544**, 389 (2002)].
- [25] A.V. Efremov, K. Goeke, and P. Schweitzer, Phys. Rev. D **67**, 114014 (2003).
- [26] F. Yuan, Phys. Lett. B **589**, 28 (2004).
- [27] L.P. Gamberg, D.S. Hwang, and K.A. Oganessian, Phys. Lett. B **584**, 276 (2004).
- [28] M. Anselmino, M. Boglione, U. D'Alesio, A. Kotzinian, F. Murgia, and A. Prokudin, Phys. Rev. D **71**, 074006 (2005).
- [29] K. Goeke, A. Metz, P.V. Pobylitsa, and M.V. Polyakov, Phys. Lett. B **567**, 27 (2003).
- [30] A. Afanasev and C.E. Carlson, arXiv:hep-ph/0308163.
- [31] A. Metz and M. Schlegel, Eur. Phys. J. **A22**, 489 (2004).

- [32] A. Bacchetta, P.J. Mulders, and F. Pijlman, Phys. Lett. B **595**, 309 (2004).
- [33] K. Goeke, A. Metz, and M. Schlegel, Phys. Lett. B **618**, 90 (2005).
- [34] J.C. Collins and D.E. Soper, Nucl. Phys. **B193**, 381 (1981) [Erratum-ibid. **B213**, 545 (1983)].
- [35] J.C. Collins, D.E. Soper, and G. Sterman, Nucl. Phys. **B250**, 199 (1985).
- [36] X. Ji, J.P. Ma, and F. Yuan, Phys. Rev. D **71**, 034005 (2005).
- [37] X. Ji, J.P. Ma, and F. Yuan, Phys. Lett. B **597**, 299 (2004).
- [38] J.C. Collins and A. Metz, Phys. Rev. Lett. **93**, 252001 (2004).
- [39] D. Boer, P.J. Mulders, and F. Pijlman, Nucl. Phys. **B667**, 201 (2003).
- [40] J.C. Collins, Acta Phys. Polon. B **34**, 3103 (2003).
- [41] J.C. Collins and D.E. Soper, Nucl. Phys. **B194**, 445 (1982).
- [42] X. Ji and F. Yuan, Phys. Lett. B **543**, 66 (2002).
- [43] D. Boer, S.J. Brodsky, D.S. Hwang, Phys. Rev. D **67**, 054003 (2003).
- [44] L.P. Gamberg, G.R. Goldstein, and K.A. Oganessyan, Phys. Rev. D **67**, 071504 (2003).
- [45] A. Bacchetta, A. Schaefer, and J.J. Yang, Phys. Lett. B **578**, 109 (2004).
- [46] K. Goeke, S. Meissner, A. Metz, and M. Schlegel, Phys. Lett. B **637**, 241 (2006).
- [47] A.V. Belitsky, X. Ji, and F. Yuan, Nucl. Phys. **B656**, 165 (2003).
- [48] J.C. Collins and F. Hautmann, Phys. Lett. B **472**, 129 (2000).
- [49] A.V. Afanasev and C.E. Carlson, arXiv:hep-ph/0603269.